

MHMH “Oral Contrast Challenge- Adult” Order Set

- Go to **Manage Orders**
- **Search for:**
 - Oral Contrast Challenge
 - Enteral contrast
 - Water soluble contrast challenge
 - Small bowel contrast challenge
 - Bedside small bowel follow through
 - Omnipaque
- **Select “Oral Contrast Challenge”**

Order and Order Set Search

ORAL CONTRAST CHALLENGE

Browse Preference List Facility List

Order Sets, Panels, & Pathways

«Search order sets by user»

P...	Name	Type
	ORAL CONTRAST CHALLENGE- ADULT	Order Panel

Medications (No results found)

- Users will be presented with five orders, one for the Omnipaque Contrast and four for the XR Abdomen
- Only the Contrast order and first XR Abdomen order have hard stop fields

Oral Contrast Challenge- Adult Accept

Order both contrast and XR imaging order when doing a contrast challenge. Timing of contrast administration and imaging need to be coordinated between nursing and radiology.

Omnipaque orders - Please select appropriate enteral route.

iohexoL (Omnipaque) (350 mg/mL) ORAL (\$\$)
100 mL, Oral, ONCE, RN to page 9217 when contrast has been administered. Radiology to schedule first x-ray 6 hours after contrast is administered., Routine

iohexoL (Omnipaque) (350 mg/mL) G TUBE (\$\$)
100 mL, Per G Tube, ONCE, RN to page 9217 when contrast has been administered. Radiology to schedule first x-ray 6 hours after contrast is administered., Routine

iohexoL (Omnipaque) (350 mg/mL) J TUBE (\$\$)
100 mL, Per J Tube, ONCE, RN to page 9217 when contrast has been administered. Radiology to schedule first x-ray 6 hours after contrast is administered., Routine

iohexoL (Omnipaque) (350 mg/mL) NG TUBE (\$\$)
100 mL, Per NG tube, ONCE, RN to page 9217 when contrast has been administered. Radiology to schedule first x-ray 6 hours after contrast is administered., Routine

XR Abdomen 1 view Contrast Challenge
Routine
To be done 6 hours after the contrast administration.
Clinical information / key questions for radiologist: first image in series

XR Abdomen 1 view Contrast Challenge
Routine
To be done 12 hours after the contrast administration.
Reason for exam and clinical history: follow up to contrast challenge
Clinical information / key questions for radiologist: second image in series

XR Abdomen 1 view Contrast Challenge
Routine
To be done 18 hours after the contrast administration.
Reason for exam and clinical history: follow up to contrast challenge
Clinical information / key questions for radiologist: third image in series

XR Abdomen 1 view Contrast Challenge
Routine
To be done 24 hours after contrast administration.
Reason for exam and clinical history: follow up to contrast challenge
Clinical information / key questions for radiologist: fourth image in series
Stat read required? No

- **Omnipaque order:**
 - Click on the specific route you would like the nurse to use to administer the contrast
 - All the necessary information is pre-filled in for you

Oral Contrast Challenge- Adult ✔ Accept

Order both contrast and XR imaging order when doing a contrast challenge. Timing of contrast administration and imaging need to be coordinated between nursing and radiology.

! Omnipaque orders - Please select appropriate enteral route.

- iohexoL (Omnipaque) (350 mg/mL) **ORAL** (\$\$)
100 mL, Oral, ONCE, RN to page 9217 when contrast has been administered. Radiology to schedule first x-ray 6 hours after contrast is administered., Routine
- iohexoL (Omnipaque) (350 mg/mL) **G TUBE** (\$\$)
100 mL, Per G Tube, ONCE, RN to page 9217 when contrast has been administered. Radiology to schedule first x-ray 6 hours after contrast is administered., Routine
- iohexoL (Omnipaque) (350 mg/mL) **J TUBE** (\$\$)
100 mL, Per J Tube, ONCE, RN to page 9217 when contrast has been administered. Radiology to schedule first x-ray 6 hours after contrast is administered., Routine
- iohexoL (Omnipaque) (350 mg/mL) **NG TUBE** (\$\$)
100 mL, Per NG tube, ONCE, RN to page 9217 when contrast has been administered. Radiology to schedule first x-ray 6 hours after contrast is administered., Routine

- **XR Abdomen orders:**
 - Enter a reason for exam
 - If the patient is female there will be an additional hardstop pregnancy question on the first order. Once an option is selected it will autofill in for the additional three orders

XR Abdomen 1 view Contrast Challenge ✔ Accept ✖ Cancel

Priority:

Time Image Should Be Taken

! Reason for exam and clinical history:

Clinical information / key questions for radiologist:

Stat read required?

- Once you are done press **Accept** and **Sign**
- This is what the medication will look like in the MAR:

Summary Chart Review Notes Manage Orders **Prep for Procedure** MAR

MAR

Report MAR Note Messages Legend Show All Actions Link Lines Anywhere RN Glu Mngmt Pain Mngmt

ALL One Time Scheduled Continuous PRN Due/Overdue Meds Override Pulls Chemo Periop Respiratory Tube Feeds Beacon

Go to Now or Select Dat... **Overdue** Show Med Education Show All Details Hide All Admins

Friday November 19, 2021

1000 | 1100 | 1200 | 1300 | **1400** | 1500 | 1600 | 1700

iohexoL (Omnipaque) (350 mg/mL) solution 100 mL Dose: 100 mL : Oral : ONCE :

1500 Due

Admin Instructions:
RN to page 9217 when contrast has been administered. Radiology to schedule first x-ray 6 hours after contrast is administered.

Ordered Admin Dose: 100 mL	Dispense Location: MHMH Radiology CONTRAST Floorstock Priority: Routine *** Barcode Scan Not Required *** **Warning Vesicant / Irritant Medication**
-------------------------------	---

****The Omnipaque does not need to be diluted and can be administered all at once****